

Review Article

Critical Appraisal of Gefitinib in the Treatment of Non-Small Cell Lung Cancer

Yung-Hung Luo^{1,2,3}, Yu-Chin Lee^{2,3}, Jacqueline Whang-Peng⁴, Yuh-Min Chen^{2,3,4*}

¹Chest Medicine Division of Medicine Department, Fenglin Branch, Taipei Veterans General Hospital, Taiwan

²Department of Chest Medicine, Taipei Veterans General Hospital, Taipei, Taiwan

³School of Medicine, National Yang-Ming University, Taipei, Taiwan

⁴Taipei Cancer Center, Taipei Medical University, Taipei, Taiwan

Abstract.

The leading cause of death from malignant tumors worldwide is lung cancer. Before the development of molecularly targeted therapy, chemotherapy with platinum-based doublets was considered as the standard first-line treatment in advanced non-small cell lung cancer (NSCLC) patients. The introduction of epidermal growth factor receptor-tyrosine kinase inhibitor (EGFR-TKI) has led to remarkable advances in the treatment of NSCLC. Two activating *EGFR* mutations (Exon 19 deletion and Exon 21 L858R mutation) have been correlated with dramatic responses to EGFR-TKI. Every effort should be made to identify the *EGFR* mutation status in NSCLC patients prior to the initial systemic treatment in order to select those who are most likely to benefit from EGFR-TKIs. At the present time, first-generation EGFR-TKIs are available for clinical use, including gefitinib and erlotinib. Gefitinib was the first drug developed as an EGFR-TKI for NSCLC treatment. Several randomized phase III studies revealed that gefitinib provided superior response rate, improved PFS, and less toxicity compared with doublet chemotherapy for advanced NSCLC with activating EGFR mutation. Currently, first-line treatment with gefitinib is used in metastatic NSCLC patients with tumor *EGFR* mutation. Gefitinib is also administered as salvage therapy for NSCLC patients previously treated with chemotherapy. The standard of care for previously untreated patients with *EGFR* mutation-negative or unknown status still remains platinum-based chemotherapy. In this article, we have reviewed the relevant clinical data regarding gefitinib as a molecularly targeted therapy for NSCLC.

Keywords : gefitinib, non-small cell lung cancer, epidermal growth factor receptor, tyrosine kinase inhibitor

綜合評論

Gefitinib 於非小細胞肺癌治療之關鍵評論

羅永鴻^{1,2,3} 李毓芹^{2,3} 彭汪嘉康⁴ 陳育民^{2,3,4*}

¹台北榮民總醫院鳳林分院 胸腔內科

²台北榮民總醫院 胸腔部

³陽明大學醫學院 內科學系

⁴台北醫學大學 台北癌症中心

中文摘要

肺癌為全世界惡性腫瘤死亡率最高之癌症。在分子標靶治療發展出來以前，以 platinum 為主的合併化學藥物治療(doublets)是晚期非小細胞肺癌的第一線標準治療。自從有了上皮細胞生長因素接收器-酪胺酸酶抑制劑(epidermal growth factor receptor-tyrosine kinase inhibitor, EGFR-TKI)之後，非小細胞肺癌的治療便有了長足之進展。*EGFR* 基因有 exon 19 突變(deletion)，或 exon 21 突變(L858R point mutation)之患者對 EGFR-TKI 的藥效反應相當地好。所以對於每一位非小細胞肺癌的患者，臨床醫師應該要盡力去檢測出患者的 *EGFR* 基因之突變狀態，以找出適合 EGFR-TKI 治療之病人。第一代的 EGFR-TKIs，包括 gefitinib 與 erlotinib。Gefitinib 是第一個發展出來治療非小細胞肺癌的 EGFR-TKI。一些隨機分配之第三期臨床試驗顯示，gefitinib 與合併化學藥物治療相比之下，對於有 *EGFR* 基因活化性突變的晚期非小細胞肺癌，具有較好的藥效反應、較好的無疾病惡化存活期、及較少的副作用。目前，gefitinib 是具有 *EGFR* 基因突變的晚期非小細胞肺癌之第一線治療藥物。Gefitinib 也可作為接受過化學治療的非小細胞肺癌患者之挽救性治療。對於野生型 *EGFR* 之患者，或者是 *EGFR* 基因突變狀態不明之患者，以 platinum 為主的合併化學藥物治療仍然是第一線標準治療。本篇文章回顧了 Gefitinib 作為分子標靶治療之相關臨床研究。

關鍵字: gefitinib、非小細胞肺癌、上皮細胞生長因素接收器、酪胺酸酶抑制劑

INTRODUCTION

Lung cancer is the leading cause of death worldwide arising from malignant tumors. Only approximately 15% of all lung cancer patients survived for 5 years or more after diagnosis. More than 85% of lung cancer cases have the non-small cell lung cancer (NSCLC) subtype; most patients present with advanced lung cancer and their 2-year survival rate is only 10-15% [1,2]. Before the development of molecularly targeted therapy, chemotherapy with platinum-based doublets was considered as the standard first-line treatment for advanced NSCLC. Chemotherapy has provided modest improvements in patient survival, but lung cancer patients generally still have a dismal prognosis. Considering the relative non-specificity and toxicity of chemotherapy, the development of novel therapeutic strategies is imperative. Over the

last decade, molecularly targeted therapies such as gefitinib or erlotinib for NSCLC treatment have demonstrated remarkable advances, including improved progression-free survival (PFS), overall survival (OS), and superior quality of life in certain patient populations [3-7].

The identification of several driver oncogenes related to tumor progression and lung cancer metastasis has resulted in the detection of several potential therapeutic targets [2,3,6]. The first oncogenic driver mutation emerged in 2004 with the detection of activating mutations in the tyrosine kinase domain of epidermal growth factor receptor (EGFR). The frequency of *EGFR* mutations in NSCLC patients has been reported to be approximately 5-30%, depending on the different population [8-11]. It was observed that a higher incidence of *EGFR* mutations was detected in East Asians [12,13]. These mutations were identified predominantly in pulmonary adenocarcinomas. EGFR and its downstream signaling pathway have been extensively studied in recent years. Two activating *EGFR* mutations (Exon 19 deletion and Exon 21 L858R point mutation) have been associated with dramatic responses to EGFR tyrosine kinase inhibitors

*Corresponding author: Yuh-Min Chen M.D., Ph.D.

*通訊作者：陳育民醫師

Tel: +886-2-28763466

Fax: +886-3-28743466

E-mail: ymchen@vghtpe.gov.tw




Figure 1. EGFR signal transduction pathway. The reactions of ligand-receptor and receptor-receptor interactions activate intracellular signaling pathways that regulate cellular survival/apoptosis, proliferation, and other cellular functions that lead to the activation of malignant behaviors. EGFR monoclonal antibody binds to EGFR and interferes with the binding of ligands (such as EGF). EGFR-TKIs (e.g. gefitinib) bind to the intracellular tyrosine kinase domain and inhibit signal transduction pathway. EGF = epidermal growth factor; EGFR = epidermal growth factor receptor; TKI = tyrosine kinase inhibitor

(TKI), such as gefitinib or erlotinib [9,10,14]. They exhibited favorable toxicity profiles and convenience for use. Gefitinib was the first drug developed as an EGFR-TKI for NCSLC treatment [15]. The relevant

clinical information regarding gefitinib as a molecularly targeted therapy for NSCLC was reviewed in this article.

SIGNAL TRANSDUCTION PATHWAY OF EGFR

Epidermal growth factor receptors (EGFR) and ErbB receptors belong to the same subclass I of the receptor tyrosine kinase (TK) superfamily. This ErbB family comprises four structurally similar receptors, including ErbB1 (EGFR/HER1), ErbB2 (HER2/Neu), ErbB3 (HER3), and ErbB4 (HER4) [16,17]. Of these four receptors, EGFR was the first to be identified. These receptors contain an extracellular domain for the binding of ligands, a transmembrane domain, and an intracellular TK domain. The binding of ligands to the extracellular domain of EGFR results in dimerization of two receptor molecules, and activates receptor autophosphorylation through TK activity. Then, autophosphorylation activates several downstream signaling pathways which may lead to tumor proliferation, migration, metastasis, and inhibition of apoptosis [16-18].

The TK family consists of two main intracellular signal transduction pathways. One of them is the Ras-Raf-MEK-MAPK pathway which regulates gene transcription, and proliferation of tumor cells. The other crucial downstream pathway is the phosphatidylinositol 3-kinase (PI3K) -Akt pathway, which activates a series of reactions including cell proliferation, growth, and blocking apoptosis. Another signaling route is through the Janus kinase/signal transducers and activators of transcription (Jak/Stat) pathway [17,18]. When all of the pathways are activated, it may ultimately contribute to various cellular reactions containing cell growth, division, repair, survival (or apoptosis), adhesion, invasion, and metastasis.

EGFR INHIBITION

In 1981, Mendelsohn and colleagues reported that EGFR is often overexpressed in the human cancer cell, and is related to a poor outcome. Thus, they proposed the theory of a targeted cancer therapy through EGFR inhibition [19]. The previous studies reported that the frequency of EGFR overexpression in NSCLC speci-

mens was around 43% to 89% depending on different populations, detection techniques, and the definition of EGFR overexpression [20-24]. However, in these studies, the association between EGFR expression and survival benefit revealed controversial results [24,25].

In these studies, several strategies were applied for targeting the EGFR. Among them, EGFR-TKIs and monoclonal antibodies (MoAbs) were most frequently used in clinical applications [1,2,4,26]. Small molecule TKIs bind to the intracellular TK domain of EGFR and interfere with signal transduction. MoAbs target the extracellular domain of EGFR and interfere with receptor signaling. EGFR-TKIs and MoAbs seem to possess similar effects with regard to receptor-dependent downstream signal transduction pathways, including the MAPK and PI3K-Akt pathways. Both of them lead to the effective inhibition of the major EGFR signaling pathways [27,28]. Therefore, EGFR-TKIs and MoAbs exhibit their anticancer effect in NSCLC patients by blocking several crucial cellular functions regulated by the EGFR, including cell-cycle arrest, cell apoptosis, blockade of cancer cell invasion and metastasis (Figure 1) [27,29,30]. After the discovery of an intimate association between the efficacy of EGFR-TKIs and activating *EGFR* mutations, the EGFR-TKIs as targeted therapies for EGFR-mutant NSCLC have been intensively studied [2-4]. Nevertheless, anti-EGFR monoclonal antibodies (e.g. cetuximab) have been studied less extensively and have demonstrated improved outcomes in merely one randomized phase III study of cetuximab added to doublet chemotherapy for treatment in chemotherapy-naïve NSCLC patients [2-4,31].

FIRST-GENERATION EGFR-TKI

First-generation EGFR-TKIs include gefitinib (Iressa) and erlotinib (Tarceva). Both have been approved for the treatment of advanced NSCLC, and are orally administered small molecules that reversibly inhibit the EGFR tyrosine kinase and interfere with downstream signaling. Their side effects (like dose-

dependent skin rash and diarrhea) are usually mild to moderate, and manageable in the outpatient department under close monitoring. The first drug developed as an EGFR-TKI for NSCLC treatment was gefitinib [2,3,15]. In the phase I trial, gefitinib has demonstrated objective responses in NSCLC patients who were heavily treated with prior extensive rounds of chemotherapy or chemoradiotherapy, when administered as a single agent [32]. Two randomized multicenter phase II studies [Iressa Dose Evaluation in Advanced Lung Cancer 1 (IDEAL 1) and 2 (IDEAL 2)] have been conducted in patients with locally advanced or metastatic NSCLC who had progressive disease after prior chemotherapy, including platinum-based and docetaxel-based therapies in order to evaluate the safety and efficacy of two oral doses of gefitinib (250 and 500 mg/day) [33,34]. In the IDEAL 1 trial, a response rate of 18.7% and remarkably improved disease-related symptoms were demonstrated. Gefitinib at 250 mg/day was as effective as 500 mg/day, and exhibited the superior toxicity profiles [33]. The IDEAL 2 trial revealed 12% objective response rates in the 250 mg/d group, and 9% in the 500 mg/d group. Similar toxicity profiles had been observed in the IDEAL 1 and IDEAL 2 studies [33,34]. Based on these trials, The US Food and Drug Administration (FDA) approved gefitinib as a third-line treatment for locally advanced or metastatic NSCLC patients after failure of platinum- and docetaxel-based chemotherapies in 2003 [18].

The ISEL (Iressa Survival Evaluation in Lung Cancer) trial requested by the US FDA was a randomized phase III study designed to evaluate the survival benefit of gefitinib [35]. In this study, 1692 patients with previously treated advanced NSCLC were randomized to either gefitinib treatment or placebo, and gefitinib did not demonstrate statistically longer survival time compared with the placebo among all patients (median survival: 5.6 vs. 5.1 months; $p = 0.087$). Nevertheless, preplanned subgroup analyses revealed that Asian patients receiving gefitinib ($n = 235$) had significantly longer survival as compared

with those receiving placebo ($n = 107$) (median survival: 9.5 vs. 5.5 months; $p = 0.01$). Significantly longer survival was also observed in patients who never smoked receiving gefitinib ($n = 250$) than those receiving a placebo ($n = 125$) (median survival: 8.9 vs. 6.1 months; $p = 0.012$). In 2005, the US FDA limited the use of gefitinib to patients continuing to derive benefit from gefitinib prescribed previously or those enrolled in the clinical trial, because the lack of survival benefit that was demonstrated in the ISEL study [15,18].

A multicenter phase III randomized trial [Iressa NSCLC Trial Evaluating Response and Survival versus Taxotere (INTEREST)] with a non-inferiority design to compare gefitinib with docetaxel recruited 1433 advanced NSCLC patients receiving at least one line of platinum-based chemotherapy. In this INTEREST trial, the results revealed the non-inferiority of gefitinib over docetaxel in terms of overall survival [median survival: 7.6 vs. 8.0 months; hazard ratio (HR): 1.020; 96% confidence interval (CI): 0.905–1.150], and suggested that gefitinib is an efficacious therapy in previously treated patients with molecularly unselected advanced NSCLC [36].

In the subgroup analyses of the ISEL study described above [35], gefitinib demonstrated significant survival benefits in East Asian patients, and those who had never smoked. Therefore, the Iressa Pan-Asia Study (IPASS) was initiated to evaluate the efficacy of gefitinib in East Asian patients [5]. The IPASS trial was the first randomized phase III trial to compare EGFR-TKI (gefitinib) with doublet chemotherapy (paclitaxel plus carboplatin) as first-line therapy in selected East Asia patients who were former light smokers or non-smokers with advanced pulmonary adenocarcinoma. In this study, 1217 patients were enrolled, and 437 patients (35.9%) received *EGFR* mutation analysis. In a subgroup including 261 cases with *EGFR* mutation positive tumor, patients who received gefitinib had significantly longer progression-free survival (PFS) than those receiving paclitaxel plus

Table 1. Phase III randomized trials comparing first-line EGFR-TKI with chemotherapy in clinically or molecularly selected NSCLC patients with activating *EGFR* mutations

	I-PASS		NEJ002		WJTOG3405		OPTIMAL (CTONG-0802)		EURTAC	
	G	P + Cb (<i>p</i> *)	G	P + Cb (<i>p</i> *)	G	D + Cis (<i>p</i> *)	E	Ge + Cb (<i>p</i> *)	E	Platinum based (<i>p</i> *)
RR (%)	71.2	47.3 (<0.001)	73.7	30.7 (<0.001)	62.1	32.2 (<0.0001)	83	36 (<0.001)	54.5	10.5
CR (%)	NA	NA	4.4	0	NA	NA	2	0	3	0
PFS (mo)	9.5	4.9 (<0.001)	10.8	5.4 (<0.001)	9.2	6.3 (<0.0001)	13.1	4.6 (<0.001)	9.7	5.2 (<0.001)
OS (mo)	21.6	21.9 (0.99)	30.5	23.6 (0.31)	34.8	37.3	22.7	28.8 (0.69)	19.3	19.5 (0.87)

*The *p* values: when comparing with EGFR-TKI group

Cb = carboplatin; Cis = cisplatin; CTONG = Chinese Thoracic Oncology Group; D = docetaxel; E = erlotinib; EGFR = epidermal growth factor receptor; EURTAC = European Tarceva versus chemotherapy; G = gefitinib; Ge = gemcitabine; I-PASS = Iressa Pan Asia study; NA = not applicable; NEJ002 = North East Japan 002; NSCLC = non-small-cell lung cancer; OPTIMAL = Erlotinib versus chemotherapy as first-line treatment for patients with advanced EGFR mutation-positive NSCLC; OS = overall survival; P = paclitaxel; PFS = progression-free survival; RR = response rate; TKI = tyrosine kinase inhibitor; WJTOG3405 = West Japan thoracic oncology group 3405

carboplatin (HR: 0.48; 95% CI: 0.36-0.64; *p* < 0.001). However, in another subgroup including 176 patients without *EGFR* mutation, PFS was significantly shorter among patients who received gefitinib than those who received paclitaxel plus carboplatin (HR: 2.85; 95% CI: 2.05-3.98; *p* < 0.001). This trial suggested that the existence of tumor *EGFR* mutations is an important predictor of better outcome in patients receiving gefitinib. The importance of identifying the *EGFR* mutation status before deciding on initial treatment was first discovered in the IPASS trial [5,37]. Similar results were further confirmed in other phase III trials that revealed a significantly improved PFS and better quality of life in patients receiving EGFR-TKI as compared with platinum-based doublet chemotherapy (Table 1) [5,38-42]. A phase III randomized trial (WJTOG 3405) of gefitinib compared with cisplatin/docetaxel as first-line treatment for EGFR-mutant NSCLC revealed that gefitinib provided a 3-year survival benefit for EGFR-mutant advanced NSCLC after 5 years of follow-up (median survival: 34.8 vs. 37.3 months; HR: 1.252; 95% CI: 0.883-1.775), although no significant difference in overall survival was observed [38,39].

In 2009, mainly based on results from IPASS and INTEREST trial, the European Medicines Agency (EMA) approved gefitinib as a targeted therapy for locally advanced or metastatic NSCLC harboring activating *EGFR* mutation [43,44]. Gefitinib is also approved for use in Japan, Taiwan, China, Korea, and

several Asian countries [18,43].

GEFITINIB COMBINED WITH CHEMOTHERAPY

Previous reports suggest that gefitinib does not induce myelosuppression, which was found in patients receiving conventional chemotherapy [17,32-35]. In order to evaluate whether the addition of EGFR-TKI to chemotherapy can enhance the antitumor effect and provide longer survival than chemotherapy alone, four phase III trials in previously untreated patients with advanced NSCLC were initiated (Table 2) [45-48]. The INTACT (Iressa NSCLC Trial Assessing Combination Treatment) 1 trial enrolled 1093 patients who were randomized to chemotherapy (gemcitabine/cisplatin) with either gefitinib 250 mg/day, gefitinib 500 mg/day, or placebo. No significantly different response rate, time to progression, and overall survival was observed among these treatment groups. The INTACT 2 study with the design similar to INTACT 1 recruited 1037 patients, and the chemotherapy regimen was carboplatin and paclitaxel [46]. The results also showed no significantly different response rate, time to progression, and overall survival among different treatment groups. The addition of Erlotinib to chemotherapy has also been evaluated, like gefitinib. In the TRIBUTE (Tarceva Responses in Conjunction with Paclitaxel and Carboplatin) and TALENT (Tarceva Lung Cancer Investigation) studies, chemotherapy-naïve NSCLC patients received chemotherapy both

Table 2. Four randomized Phase III trials of chemotherapy with or without EGFR-TKI (gefitinib or erlotinib) in clinical or molecularly-unselected patients with treatment-naïve advanced NSCLC (A total of 4361 patients)

Trial	No. of patients	Treatment comparison	Response rate (%)	PFS (mo)	OS (mo)
INTACT 1	1093	Cis + gemcitabine, or with gefitinib (250 or 500 mg/d)	44.8 vs. 50.3 vs. 49.7	6.0 vs. 5.8 vs. 5.5 (<i>p</i> = 0.76)	10.9 vs. 9.9 vs. 9.9 (<i>p</i> = 0.45)
INTACT 2	1037	Cb + paclitaxel, or with gefitinib (250 or 500 mg/d)	42 vs. 41 vs. 37	5.0 vs. 5.3 vs. 4.6 (<i>p</i> = 0.562)	9.9 vs. 9.8 vs. 8.7 (<i>p</i> = 0.64)
TRIBUTE	1059	Cb + paclitaxel, or with erlotinib	19.3 vs. 21.5	4.9 vs. 5.1 (<i>p</i> = 0.36)	10.5 vs. 10.6 (<i>p</i> = 0.95)
TALENT	1172	Cis + gemcitabine, or with erlotinib	29.9 vs. 31.5	5.7 vs. 5.5 (<i>p</i> = 0.74)	10.3 vs. 10.0 (<i>p</i> = 0.49)

Cb = carboplatin; Cis = cisplatin; INTACT = Iressa NSCLC Trial Assessing Combination Treatment; NSCLC = non-small-cell lung cancer; PFS = progression-free survival; OS = overall survival; TALENT = Tarceva Lung Cancer Investigation; TKI = tyrosine kinase inhibitor; TRIBUTE = Tarceva Responses in Conjunction with Paclitaxel and Carboplatin

with and without erlotinib [47,48]. These trials also failed to demonstrate improved survival in patients receiving combination therapy with chemotherapy plus erlotinib. The above-mentioned four randomized phase III studies, including 4361 patients with chemo-naïve NSCLC, obviously revealed that the combination of EGFR-TKIs (gefitinib or erlotinib) with chemotherapy does not provide superior survival benefit over chemotherapy alone. Therefore, gefitinib should not be added to cytotoxic chemotherapy as the first-line treatment in molecularly unselected NSCLC patients [2,3].

EGFR MUTATION

Previous studies have demonstrated that *EGFR* mutations may predict treatment response to EGFR-TKIs and also be a prognostic factor [17,47,49–53]. Cappuzzo and colleagues analyzed tumor specimens of NSCLC from 102 patients receiving gefitinib for tumor *EGFR* status by fluorescence *in-situ* hybridization (FISH), and immunohistochemistry, and for *EGFR* mutation by DNA sequencing [49]. The results revealed the tumor *EGFR* status and mutation were correlated with treatment response and patient survival [2,3,49]. *EGFR* mutation status possessed the best correlation with treatment effect [3,9,10,17,54], followed by FISH assay [49,54]. Several large serial studies on the epidemiology of *EGFR* mutation revealed that the reported mutations were all somatic and identified in

exons 18, 19, 20, and 21 of the *EGFR* gene encoding partial intracellular TK domain of EGFR [13,55,56]. Prior studies also reported that the sensitivity to EGFR-TKIs was correlated with three kinds of *EGFR* mutations, including (1) in-frame deletions within exon 19, accounting for approximately 46% of all EGFR mutations (the most common EGFR mutations); (2) missense mutations within exon 18, 20, or 21, accounting for approximately 41% of all mutations (the second most common EGFR mutations), especially L858R point mutations within exon 21; and (3) in-frame duplications and/or insertions within exon 20 (approximately 5% of all mutations) [2,13,55]. In addition to *EGFR* mutations, East Asian descent, female sex, never-smoker, and adenocarcinoma histology have also been reported to have a correlation with sensitivity to EGFR-TKIs. Another large study that enrolled 12,244 patients revealed that the most common mutations were exon 19 mutation (50%), followed by exon 21 (40%), exon 20 (6%), and exon 18 (4%) mutations. L858R in exon 21 and deletion of E746-A750 in exon 19 accounted for approximately 33% and 24% of all mutations [56]. An obvious correlation between the response to EGFR-TKIs and EGFR mutations has been reported. The best response rate (around 70%) was demonstrated in NSCLC patients with exon 19 mutations, followed by those with exon 21, 18, and 20 mutations (approximately 20% or slightly higher). EGFR-mutant NSCLC patients receiving EGFR-TKI

Table 3. Six phase III trials of maintenance therapy with EGFR-TKIs after first line chemotherapy (A total of 3006 patients)

Trial	No. of patients	Maintenance treatment	Median PFS	<i>p</i> -value	Median OS (mo)	<i>p</i> -value
WJTOG 0203	595	Gefitinib vs. observation	4.6 vs. 4.3 months	<0.001	13.7 vs. 12.9	0.11
EORTC 08021-ILCP 01/03	173	Gefitinib vs. placebo	4.1 vs. 2.9 months	0.002	10.9 vs. 9.4	0.204
INFORM	296	Gefitinib vs. placebo	4.8 vs. 2.6 months	<0.0001	18.7 vs. 16.9	0.26
SATURN	889	Erlotinib vs. placebo	12.3 vs. 11.1 weeks	<0.0001	12 vs. 11	0.0088
ATLAS	743	Bevacizumab + erlotinib vs. Bevacizumab + placebo	4.8 vs. 3.7 months	0.001	14.4 vs. 13.3	0.5604
IFCT-GFPC 0502 ^a	310	Erlotinib vs. observation	2.9 vs. 1.9 months	0.003	11.4 vs. 10.8	0.3043

Abbreviations: ATLAS = Avastin Tarceva Lung Adenocarcinoma Study; EGFR = epidermal growth factor receptor; EORTC = European Organization for Research and Treatment of Cancer; IFCT-GFPC = Intergroupe Francophone de Cancerologie Thoracique–Groupe Francais de Pneumo-Cancerologie; ILCP = Italian Lung Cancer Project; OS = overall survival; PFS = progression-free survival; TKI = tyrosine kinase inhibitor; SATURN = Sequential Tarceva in Unresectable Non–Small-Cell Lung Cancer; WJTOG = West Japan Thoracic Oncology Group

^aIFCT-GFPC 0502 was a trial with 3 arms, including maintenance gemcitabine, maintenance erlotinib, and observation group

had median PFS of about 10 months. The PFS varied widely because of several potential mechanisms, including T790M mutation, BIM polymorphism, c-Met amplification, and alterations within chromosome 7p, and even the percentage of *EGFR* mutation-negative cancer cells [2–4,57–62].

Several methods are utilized to identify *EGFR* mutations in tumor specimens from lung cancer patients, including direct sequencing, Scorpion ARMS (amplified refractory mutation system), and PNA-LNA (peptide nucleic acid-locked nucleic acid) PCR (polymerase chain reaction) clamp techniques [5, 63–66]. The sensitivity of direct sequencing is relatively low and it needs a higher percentage of mutant DNA in order to detect *EGFR* mutations in tumor specimens (10–25% mutant DNA). Therefore, objective response rates of 10–20% or higher in particular clinical populations receiving EGFR-TKI were reported in *EGFR* wild type NSCLC patients whose *EGFR* mutation status was detected by the direct DNA sequencing method [67–69]. High-sensitivity methods, including Scorpion ARMS and PNA-LNA PCR clamp techniques, are capable of detecting $\geq 1\%$ mutant DNA [5,63–65]. However, the Scorpion ARMS and PNA-LNA PCR clamp techniques are only able to identify recognized mutations, but direct sequencing is capable of detecting new mutations [63–66]. Tumor specimens used for *EGFR* mutation assessment are

commonly collected from biopsies in the detected pulmonary or metastatic tumors. Specimens obtained from pleural effusion, plasma-free DNA, and circulating tumor cells may also be utilized for *EGFR* mutation assessment [63,64,66,70–73].

FIRST-LINE GEFITINIB TREATMENT FOR EGFR-MUTANT NSCLC

First-line treatment with EGFR-TKIs for EGFR-mutant NSCLC has demonstrated improved PFS, better quality of life, and less treatment-related toxicity compared with platinum-based doublet chemotherapy in several recent randomized phase III studies (Table 1) [5,38–42]. Thus, *EGFR* mutation analysis for advanced NSCLC patients being considered for first-line EGFR-TKI should be performed, and gefitinib as first-line treatment is recommended for NSCLC patients with activating *EGFR* mutation [74]. EGFR-mutant NSCLC patients receiving EGFR-TKI had median PFS of around 10 months. When first-line platinum-based doublet chemotherapy was used for EGFR-mutant NSCLC, the median PFS was only 5–6 months. However, no significant difference in overall survival was found between EGFR-mutant patients receiving EGFR-TKI and those receiving first-line chemotherapy [41,75–78]. In patients with *EGFR* mutation-negative or unknown status, first-line treatment

with chemotherapy is preferred. First-line EGFR-TKIs in NSCLC patients with *EGFR* mutation-negative or unknown status might have a detrimental effect on PFS and OS [5,79,80].

In spite of the initial successful treatment with gefitinib in EGFR-mutant NSCLC patients, most of them will eventually experience progressive disease (acquired resistance to gefitinib) [81]. Strategies for overcoming acquired resistance to gefitinib are being developed [2,3,57]. At the present time, a number of treatment choices are available after failure of gefitinib treatment, such as salvage chemotherapy, use of another approved EGFR-TKI, participating in clinical studies with second-line molecularly targeted therapy, or chemotherapy with continuation of the EGFR-TKI already initiated. Of the above-mentioned therapeutic choices, only salvage chemotherapy is regarded as a standard of care or a preferred option after failure of gefitinib [2-4].

GEFITINIB AS MAINTENANCE TREATMENT AFTER FIRST-LINE CHEMOTHERAPY

The strategy of maintenance therapy with EGFR-TKIs after first-line treatment has been assessed in several trials. Two phase III trials with molecularly unselected patients have demonstrated that gefitinib was not indicated as a maintenance or an adjuvant therapy in stage III patients after chemoradiotherapy or in patients with completely resected NSCLC [82,83]. In a retrospective study with resected stage I to III pulmonary adenocarcinoma patients harboring *EGFR* exon 19 deletions or exon 21 L858R mutation, a tendency toward improved disease-free survival was discovered in patients receiving adjuvant treatment with gefitinib or erlotinib compared with those who did not receive similar treatment [84]. Nevertheless, additional investigation is warranted to verify the above-mentioned results.

A number of phase III trials in clinically or molecularly unselected patients with advanced NSCLC have

also revealed that gefitinib was not indicated for combined treatment with EGFR-TKI and chemotherapy followed by a maintenance therapy with EGFR-TKI alone (Table 2) [45-48]. However, whether an intercalated combination of EGFR-TKI and chemotherapy provides clinical benefit remains debatable [85-87]. Switch maintenance therapy with EGFR-TKIs after first-line chemotherapy has been evaluated in 6 phase III trials (Table 3) [88-94]. In the West Japan Thoracic Oncology Group trial (WJTOG0203), advanced NSCLC patients were randomized to 6 cycles of doublet chemotherapy or 3 cycles of chemotherapy followed by gefitinib until the disease progressed [88]. Significantly improved PFS was observed in the patients receiving chemotherapy followed by gefitinib compared with those receiving chemotherapy alone, although no significant difference in OS was demonstrated. Erlotinib has also been evaluated as a maintenance treatment in the Sequential Tarceva in Unresectable NSCLC (SATURN) trial including 889 molecularly unselected NSCLC patients without progressive disease after doublet chemotherapy for 4 cycles [91]. Erlotinib maintenance therapy yielded significant improvement in PFS (HR: 0.71; $p < 0.0001$), and overall survival compared with placebo (median survival: 12 vs. 11 months; $p = 0.0088$). Similar results were demonstrated in the other studies [92-94].

Similar to the SATURN study, the Iressa in NSCLC for maintenance (INFORM) trial revealed that the gefitinib maintenance therapy provided longer median PFS than placebo (4.8 vs. 2.6 months; $p < 0.0001$), but no significant difference in OS was observed. The PFS of patients receiving gefitinib maintenance therapy was significantly improved in the EGFR-mutant patients (HR: 0.17; 95% CI: 0.07-0.42), but not in EGFR mutation-negative patients (HR: 0.86, 95% CI: 0.48-1.51). In patients with squamous cell carcinoma, no improvement in PFS was also found. The gefitinib maintenance treatment demonstrated a greater benefit in patients with activating *EGFR* mutation [88,91,95].

ADVERSE EFFECTS OF GEFITINIB

The main adverse reactions of gefitinib are dermatologic and gastrointestinal toxicities, including skin rash/acneiform rashes (29-66%), dry skin (15-31%), pruritus (19-45%), paronychia (1-14%), diarrhea (16-47%), and elevated aminotransferase (11-26%) [5,33,35,36,42,76]. Most toxicities were usually mild to moderate. In the IPASS trial, neutropenia (3.7%) and anemia (2.2%) have been reported [5]. Interstitial lung disease (ILD) related to gefitinib has also been observed. An analysis of 50,005 patients initiated by the US FDA revealed that the worldwide incidence of ILD related to gefitinib treatment was approximately 1% [higher in Japan (2%) than in USA (0.3%)], and about one-third of these cases were lethal [96,97]. In a number of recent phase III trial, the reported incidence of ILD ranged from 1.3 to 5.3 % [5,38,40,76]. Gefitinib treatment clearly demonstrated a favorable toxicity profile compared with doublet chemotherapy.

CONCLUSIONS

Gefitinib provides better response rate, improved PFS, and less toxicity compared with platinum-based doublet chemotherapy for advanced NSCLC with activating *EGFR* mutation. Every effort should be made to identify the *EGFR* mutation status before the initial systemic therapy for NSCLC patients. First-line treatment with EGFR-TKI, including gefitinib or erlotinib, should be used in patients with EGFR-mutant advanced NSCLC. For patients with unknown *EGFR* mutation status and poor performance status (PS 3-4), standard chemotherapy is an inappropriate treatment, so EGFR-TKI as first-line therapy could be considered for them. EGFR-mutant NSCLC patients receiving gefitinib had median PFS of approximately 10 months. Further investigations about gefitinib are mandatory in order to develop better treatment strategies and optimal therapy for each patient.

REFERENCES

1. Ettinger DS, Akerley W, Borghaei H, et al. Non-small cell lung cancer. **J Natl Compr Canc Netw** **10**: 1236-71, 2012.
2. Chen YM. Update of epidermal growth factor receptor-tyrosine kinase inhibitors in non-small-cell lung cancer. **J Chin Med Assoc** **76**: 249-57, 2013.
3. Chen YM, Whang-Peng J, Chen CM. Review of first-line systemic therapy for metastatic non-small-cell lung cancer. **J Exp Clin Med** **3**: 116-20, 2011.
4. Azzoli CG, Baker Jr S, Temin S, et al. American Society of Clinical Oncology Clinical Practice Guideline update on chemotherapy for stage IV non-small-cell lung cancer. **J Clin Oncol** **27**: 6251-66, 2009.
5. Mok TS, Wu YL, Thongprasert S, et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. **N Engl J Med** **361**: 947-57, 2009.
6. Shepherd FA, Rodrigues PJ, Ciuleanu T, et al. Erlotinib in previously treated non-small-cell lung cancer. **N Engl J Med** **353**: 123-32, 2005.
7. Thongprasert S, Duffield E, Saijo N, et al. Health-related quality-of-life in a randomized phase III first-line study of gefitinib versus carboplatin/paclitaxel in clinically selected patients from Asia with advanced NSCLC (IPASS). **J Thorac Oncol** **6**: 1872-80, 2011.
8. Bronte G, Rizzo S, La Paglia L, et al. Driver mutations and differential sensitivity to targeted therapies: a new approach to the treatment of lung adenocarcinoma. **Cancer Treat Rev** **36**: S21-9, 2010.
9. Lynch TJ, Bell DW, Sordell R, et al. Activating mutations in the epidermal growth factor receptor underlying responsiveness of non-small-cell lung cancer to gefitinib. **N Engl J Med** **350**: 2129-39, 2004.
10. Paez JG, Janne PA, Lee JC, et al. EGFR mutations in lung cancer: correlation with clinical response

- to gefitinib therapy. **Science** **304**: 1497-500, 2004.
11. Inoue A, Suzuki T, Fukuhara T, et al. Prospective phase II study of gefitinib for chemotherapy-naïve patients with advanced non-small-cell lung cancer with epidermal growth factor receptor gene mutations. **J Clin Oncol** **24**: 3340-6, 2006.
 12. Kosaka T, Yatabe Y, Endoh H, et al. Mutations of the epidermal growth factor receptor gene in lung cancer: biological and clinical implications. **Cancer Res** **64**: 8919-23, 2004.
 13. Shigematsu H, Lin L, Takahashi T, et al. Clinical and biological features associated with epidermal growth factor receptor gene mutations in lung cancers. **J Natl Cancer Inst** **97**: 339-46, 2005.
 14. Pao W, Miller V, Zakowski M, et al. EGF receptor gene mutations are common in lung cancers from “never smokers” and are associated with sensitivity of tumors to gefitinib and erlotinib. **Proc Natl Acad Sci USA** **101**: 13306-11, 2004.
 15. Mahipal A, Kothari N, Gupta S. Epidermal growth factor receptor inhibitors: coming of age. **Cancer Control** **21**: 74-9, 2014.
 16. Hodgkinson PS, MacKinnon A, Sethi T. Targeting growth factors in lung cancer. **Chest** **133**: 1209-16, 2008.
 17. Chen YM. Clinical application of epidermal growth factor receptor tyrosine kinase inhibitors (EGFR-TKI) against non-small cell lung cancer (NSCLC). **J Cancer Molecules** **1**: 83-91, 2005.
 18. Ciardiello F, Tortora G. EGFR antagonists in cancer treatment. **N Engl J Med** **358**: 1160-74, 2008.
 19. Mendelsohn J. Targeting the epidermal growth factor receptor for cancer therapy. **J Clin Oncol** **20**: 1S-13S, 2002.
 20. Pastorino U, Andreola S, Tagliabue E, et al. Immunocytochemical markers in stage I lung cancer: relevance to prognosis. **J Clin Oncol** **15**: 2858-65, 1997.
 21. Greatens TM, Niehans GA, Rubins JB, et al. Do molecular markers predict survival in non-small-cell lung cancer? **Am J Respir Crit Care Med** **157**: 1093-7, 1998.
 22. Fu XL, Zhu XZ, Shi DR, et al. Study of prognostic predictors for non small cell lung cancer. **Lung Cancer** **23**: 143-52, 1999.
 23. Ohsaki Y, Tanno S, Fujita Y, et al. Epidermal growth factor receptor expression correlates with poor prognosis in non-small cell lung cancer patients with p53 over expression. **Oncol Rep** **7**: 603-7, 2000.
 24. Hirsch FR, Varella-Garcia M, Bunn PA Jr, et al. Epidermal growth factor receptor in non-small-cell lung carcinomas. Correlation between gene copy number and protein expression and impact on prognosis. **J Clin Oncol** **21**: 3798-807, 2003.
 25. Meert AP, Martin B, Delmotte P, et al. The role of EGFR expression on patient survival in lung cancer: a systemic review with meta-analysis. **Eur Respir J** **20**: 975-81, 2002.
 26. Baselga J, Arteaga CL. Critical update and emerging trends in epidermal growth factor receptor targeting in cancer. **J Clin Oncol** **23**: 2445-59, 2005.
 27. Mendelsohn J, Baselga J. Status of epidermal growth factor receptor antagonists in the biology and treatment of cancer. **J Clin Oncol** **21**: 2787-99, 2003.
 28. Pao W, Miller VA. Epidermal growth factor receptor mutations, small molecules kinase inhibitors, and non-small-cell lung cancer: current knowledge and future directions. **J Clin Oncol** **23**: 2556-68, 2005.
 29. Woodburn JR. The epidermal growth factor receptor and its inhibition in cancer therapy. **Pharmacol Ther** **82**: 241-50, 1999.
 30. Mendelsohn J, Baselga J. The EGF receptor family as targets for cancer therapy. **Oncogene** **19**: 6550-65, 2000.
 31. Pirker R, Pereira JR, Szczesna A, et al. Cetuximab plus chemotherapy in patients with advanced non-small cell lung cancer (FLEX): an open-label randomised phase III trial. **Lancet** **373**: 1525-31,

- 2009.
32. Ranson M, Hammond LA, Ferry D, et al. ZD1839, a selective oral epidermal growth factor receptor-tyrosine kinase inhibitor, is well tolerated and active in patients with solid, malignant tumors: results of a phase I trial. **J Clin Oncol** **20**: 2240-50, 2002.
 33. Fukuoka M, Yano S, Giaccone G, et al. Multi-institutional randomized phase II trial of gefitinib for previously treated patients with advanced non-small-cell lung cancer (the IDEAL 1 trial). **J Clin Oncol** **21**: 2237-46, 2003.
 34. Kris MG, Natale RB, Herbst RS, et al. Efficacy of gefitinib, an inhibitor of the epidermal growth factor receptor tyrosine kinase, in symptomatic patients with non-small cell lung cancer: a randomized trial. **JAMA** **290**: 2149-58, 2003.
 35. Thatcher N, Chang A, Parikh P, et al. Gefitinib plus best supportive care in previously treated patients with refractory advanced non-small-cell lung cancer: results from a randomised, placebo-controlled, multicentre study (Iressa Survival Evaluation in Lung Cancer). **Lancet** **366**: 1527-37, 2005.
 36. Kim ES, Hirsh V, Mok T, et al. Gefitinib versus docetaxel in previously treated non-small-cell lung cancer (INTEREST): a randomised phase III trial. **Lancet** **372**: 1809-18, 2008.
 37. Jett JR, Carr LL. Targeted therapy for non-small cell lung cancer. **Am J Respir Crit Care Med** **188**: 907-12, 2013.
 38. Mitsudomi T, Morita S, Yatabe Y, et al. Gefitinib versus cisplatin plus docetaxel in patients with non-small-cell lung cancer harbouring mutations of the epidermal growth factor receptor (WJTOG3405): an open label, randomised phase 3 trial. **Lancet Oncol** **11**: 121-8, 2010.
 39. Yoshioka H, Mitsudomi T, Morita S, et al. Final overall survival results of WJTOG 3405, a randomized phase 3 trial comparing gefitinib (G) with cisplatin plus docetaxel (CD) as the first-line treatment for patients with non-small cell lung cancer (NSCLC) harboring mutations of the epidermal growth factor receptor (EGFR). **J Clin Oncol** **32** (5 suppl): abstract 8117, 2014.
 40. Maemondo M, Inoue A, Kobayashi K, et al. Gefitinib or chemotherapy for non-small-cell lung cancer with mutated EGFR. **N Engl J Med** **362**: 2380-8, 2010.
 41. Zhou C, Wu YL, Chen G, et al. Erlotinib versus chemotherapy as first-line treatment for patients with advanced EGFR mutation-positive non-small-cell lung cancer (OPTIMAL, CTONG-0802): a multicentre, open-label, randomised, phase 3 study. **Lancet Oncol** **12**: 735-42, 2011.
 42. Rosell R, Carcereny E, Gervais R, et al. Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EURTAC): a multicentre, open-label, randomised phase 3 trial. **Lancet Oncol** **13**: 239-46, 2012.
 43. Gridelli C, De Marinis F, Di Maio M, et al. Gefitinib as first-line treatment for patients with advanced non-small-cell lung cancer with activating epidermal growth factor receptor mutation: Review of the evidence. **Lung Cancer** **71**: 249-57, 2011.
 44. Köhler JI, Schuler M. Afatinib, erlotinib and gefitinib in the first-line therapy of EGFR mutation-positive lung adenocarcinoma: a review. **Onkologie** **36**: 510-8, 2013.
 45. Giaccone G, Herbst RS, Manegold C, et al. Gefitinib in combination with gemcitabine and cisplatin in advanced non-small-cell lung cancer: a phase III trial—INTACT 1. **J Clin Oncol** **22**: 777-84, 2004.
 46. Herbst RS, Giaccone G, Schiller JH, et al. Gefitinib (ZD1839) in combination with paclitaxel and carboplatin in advanced non-small-cell lung cancer: a phase III trial—INTACT 2. **J Clin Oncol** **22**: 785-94, 2004.
 47. Herbst RS, Prager D, Hermann R, et al. TRIB-

- UTE: a phase III trial of erlotinib hydrochloride (osi-774) combined with carboplatin and paclitaxel chemotherapy in advanced non-small-cell lung cancer. **J Clin Oncol** **23**: 5892-9, 2005.
48. Gatzemeier U, Pluzanska A, Szczesna A, et al. Phase III study of erlotinib in combination with cisplatin and gemcitabine in advanced non-small cell lung cancer: the Tarceva Lung Cancer Investigation Trial. **J Clin Oncol** **25**: 1545-52, 2007.
 49. Cappuzzo F, Hirsch FR, Rossi E, et al. Epidermal growth factor receptor gene and protein and gefitinib sensitivity in non-small-cell lung cancer. **J Natl Cancer Inst** **97**: 643-5, 2005.
 50. Han SW, Kim TY, Hwang PG, et al. Predictive and prognostic impact of epidermal growth factor receptor mutation in non-small-cell lung cancer patients treated with gefitinib. **J Clin Oncol** **23**: 2493-501, 2005.
 51. Mitsudomi T, Kosaka T, Endoh H, et al. Mutations of the epidermal growth factor receptor gene predict prolonged survival after gefitinib treatment in patients with non-small-cell lung cancer with postoperative recurrence. **J Clin Oncol** **23**: 2513-20, 2005.
 52. Luo YH, Wu CH, Wu WS, et al. Association between tumor epidermal growth factor receptor mutation and pulmonary tuberculosis in patients with adenocarcinoma of the lungs. **J Thorac Oncol** **7**: 299-30, 2012.
 53. Luo YH, Ho HL, Tsai CM, et al. The Association between tumor epidermal growth factor receptor (EGFR) mutation and multiple primary malignancies in patients with adenocarcinoma of the lungs. **Am J Clin Oncol** **38**: 147-51, 2015.
 54. Hirsch FR, Dziadziuszko R, Varella-Garcia M, et al. First-generation epidermal growth factor receptor inhibitors in non-small cell lung cancer: clinical impact of the epidermal growth factor receptor fluorescence in situ hybridization assay. **J Thorac Oncol** **3**: S138-42, 2008.
 55. Shigematsu H, Gazdar A. The epidemiology of EGFR mutations. **Signal** **6**: 4-8, 2005.
 56. Murray S, Dahabreh IJ, Linardou H, et al. Somatic mutations of the tyrosine kinase domain of epidermal growth factor receptor and tyrosine kinase inhibitor response to TKIs in non-small cell lung cancer: an analytical database. **J Thorac Oncol** **3**: 832-9, 2008.
 57. Su KY, Chen HY, Li KC, et al. Pretreatment epidermal growth factor receptor (EGFR) T790M mutation predicts shorter EGFR tyrosine kinase inhibitor response duration in patients with non-small-cell lung cancer. **J Clin Oncol** **30**: 433-40, 2012.
 58. Yuan S, Yu SL, Chen HY, et al. Clustered genomic alterations in chromosome 7p dictate outcomes and targeted treatment responses of lung adenocarcinoma with EGFR-activating mutations. **J Clin Oncol** **29**: 3435-42, 2011.
 59. Rosell R, Moran T, Queralt C, et al. Screening for epidermal growth factor receptor mutations in lung cancer. **N Engl J Med** **361**: 958-67, 2009.
 60. Fan WC, Yu CJ, Tsai CM, et al. Different efficacies of erlotinib and gefitinib in Taiwanese patients with advanced non-small cell lung cancer: a retrospective multi-center study. **J Thorac Oncol** **6**: 148-55, 2011.
 61. Zhou Q, Zhang XC, Chen ZH, et al. Relative abundance of EGFR mutations predicts benefit from gefitinib treatment for advanced non-small-cell lung cancer. **J Clin Oncol** **29**: 3316-21, 2011.
 62. Ng KP, Hillmer AM, Chuah CT, et al. A common BIM deletion polymorphism mediates intrinsic resistance and inferior responses to tyrosine kinase inhibitors in cancer. **Nat Med** **18**: 521-8, 2012.
 63. Pao W, Ladanyi M. Epidermal growth factor receptor mutation testing in lung cancer: searching for the ideal method. **Clin Cancer Res** **13**: 4954-5, 2007.
 64. Mack PC, Holland WS, Burich RA, et al. EGFR mutations detected in plasma are associated with

- patient outcomes in erlotinib plus docetaxel-treated non-small cell lung cancer. **J Thorac Oncol** 4: 1466-72, 2009.
65. Tanaka T, Nagai Y, Miyazawa H, et al. Reliability of the peptide nucleic acid-locked nucleic acid polymerase chain reaction clamp-based test for epidermal growth factor receptor mutations integrated into the clinical practice for non-small cell lung cancers. **Cancer Sci** 98: 246-52, 2007.
66. Chen YM, Fan WC, Tseng PC, et al. Plasma epidermal growth factor receptor mutation analysis and possible clinical application in pulmonary adenocarcinoma patients treated with erlotinib. **Oncol Lett** 3: 713-7, 2012.
67. Wu WS, Chen YM, Tsai CM, et al. Erlotinib has better efficacy than gefitinib in adenocarcinoma patients without tumor EGFR-activating mutation, but similar efficacy in tumor with EGFR-activating mutations. **Exp Ther Med** 3: 207-13, 2012.
68. Chen YM, Tsai CM, Fan WC, et al. Phase II randomized trial of erlotinib or vinorelbine in chemo-naïve, advanced, non-small-cell lung cancer (NSCLC) patients aged 70 years or older. **J Thorac Oncol** 7: 412-8, 2012.
69. Chen YM, Fan WC, Tsai CM, et al. A phase II randomized trial of gefitinib alone or with tegafur/uracil treatment in patients with pulmonary adenocarcinoma who had failed previous chemotherapy. **J Thorac Oncol** 6: 1110-6, 2011.
70. Goto K, Ichinose Y, Ohe Y, et al. Epidermal growth factor receptor mutation status in circulating free DNA in serum from IPASS, a phase III study of gefitinib or carboplatin/paclitaxel in non-small cell lung cancer. **J Thorac Oncol** 7: 115-21, 2012.
71. Kimura H, Kasahara K, Kawaishi M, et al. Detection of epidermal growth factor receptor mutations in serum as a predictor of the response to gefitinib in patients with non-small-cell lung cancer. **Clin Cancer Res** 12: 3915-21, 2006.
72. Tsai TH, Wu SG, Chang YL, et al. Effusion immunocytochemistry as an alternative approach for the selection of first line targeted therapy in advanced lung adenocarcinoma. **J Thorac Oncol** 7: 993-1000, 2012.
73. Nakamura T, Sueoka-Aragane N, Iwanaga K, et al. A noninvasive system for monitoring resistance to epidermal growth factor receptor tyrosine kinase inhibitors with plasma DNA. **J Thorac Oncol** 6: 1639-48, 2011.
74. Keedy VL, Temin S, Somerfield MR, et al. American Society of Clinical Oncology provisional clinical opinion: epidermal growth factor receptor (EGFR) mutation testing for patients with advanced non-small-cell lung cancer considering first-line EGFR tyrosine kinase inhibitor therapy. **J Clin Oncol** 29: 2121-7, 2011.
75. Fukuoka M, Wu YL, Thongprasert S, et al. Biomarker analyses and final overall survival results from a phase III, randomized, open-label, first-line study of gefitinib versus carboplatin/paclitaxel in clinically selected patients with advanced non-small-cell lung cancer in Asia (IPASS). **J Clin Oncol** 29: 2866-74, 2011.
76. Han JY, Park K, Kim SW, et al. First-SIGNAL: first-line single-agent Iressa versus gemcitabine and cisplatin trial in never-smokers with adenocarcinoma of the lung. **J Clin Oncol** 30: 1122-8, 2012.
77. Zhou C, Wu YL, Liu X, et al. Overall survival(OS) results from OPTIMAL (CTONG0802), a phase III trial of erlotinib (E) versus carboplatin plus gemcitabine (GC) as first-line treatment for Chinese patients with EGFR mutation-positive advanced non small cell lung cancer (NSCLC). **J Clin Oncol** 30 (15 Suppl): abstract 7520, 2012.
78. Mitsudomi T, Morita S, Yatabe Y, et al. Updated overall survival results of WJTOG 3405, a randomized phase III trial comparing gefitinib (G) with cisplatin plus docetaxel (CD) as the first-line treatment for patients with non-small cell lung cancer harboring mutations of the epidermal

- growth factor receptor (EGFR). **J Clin Oncol** **30** (15 Suppl): abstract 7521, 2012.
79. Gridelli C, Butts C, Ciardiello F, et al. An international, multicenter, randomized phase III study of first-line erlotinib followed by second-line cisplatin/gemcitabine versus first-line cisplatin/gemcitabine followed by second-line erlotinib in advanced non-small-cell lung cancer: treatment Rationale and protocol dynamics of the TORCH trial. **Clin Lung Cancer** **9**: 235-8, 2008.
 80. Thomas M, Reuss A, Fischer JR, et al. Innovations: Randomized phase II trial of erlotinib (E)/bevacizumab(B) compared with cisplatin (P)/gemcitabine(G) plus B in first-line treatment of advanced nonsquamous (NS) non-small cell lung cancer (NSCLC). **J Clin Oncol** **29** (15 suppl): abstract 7504, 2011.
 81. Jackman D, Pao W, Riely GJ, et al. Clinical definition of acquired resistance to epidermal growth factor receptor tyrosine kinase inhibitors in non-small-cell lung cancer. **J Clin Oncol** **28**: 357-60, 2010.
 82. Kelly K, Chansky K, Gaspar LE, et al. Phase III trial of maintenance gefitinib or placebo after concurrent chemoradiotherapy and docetaxel consolidation in inoperable stage III non-small-cell lung cancer: SWOG S0023. **J Clin Oncol** **26**: 2450-56, 2008.
 83. Goss GD, O'Callaghan C, Lorimer I, et al. Gefitinib versus placebo in completely resected non-small-cell lung cancer: results of the NCIC CTG BR19 study. **J Clin Oncol** **31**: 3320-26, 2013.
 84. Janjigian YY, Park BJ, Zakowski MF, et al. Impact on disease-free survival of adjuvant erlotinib or gefitinib in patients with resected lung adenocarcinomas that harbor EGFR mutations. **J Thorac Oncol** **6**: 569-75, 2011.
 85. Mok T, Wu YL, Yu CJ, et al. Randomized, placebo-controlled, phase II study of sequential erlotinib and chemotherapy as first-line treatment for advanced non-small-cell lung cancer. **J Clin Oncol** **27**: 5080-7, 2009.
 86. Hirsch FR, Kabbinar F, Eisen T, et al. A randomized, phase II, biomarker-selected study comparing erlotinib to erlotinib intercalated with chemotherapy in first line therapy for advanced non-small-cell lung cancer. **J Clin Oncol** **29**: 3567-73, 2011.
 87. Mok T, Wu YL, Thongprasert S, et al. A randomized placebo-controlled phase 3 study of intercalated erlotinib with gemcitabine/platinum in 1st-line advanced non-small-cell lung cancer (NSCLC): FASTACT-2. **J Clin Oncol** **30** (15 suppl): abstract 7519, 2012.
 88. Takeda K, Hida T, Sato T, et al. Randomized phase III trial of platinum-doublet chemotherapy followed by gefitinib compared with continued platinum-doublet chemotherapy in Japanese patients with advanced non-small-cell lung cancer: results of a west Japan thoracic oncology group trial (WJTOG0203). **J Clin Oncol** **28**: 753-60, 2010.
 89. Gaafar RM, Surmont VF, Scagliotti GV, et al. A double-blind, randomised, placebo-controlled phase III intergroup study of gefitinib in patients with advanced NSCLC, non-progressing after first line platinum-based chemotherapy (EORTC 08021/ILCP01/03). **Eur J Cancer** **47**: 2331-40, 2011.
 90. Zhang L, Ma S, Song X, et al. Gefitinib versus placebo as maintenance therapy in patients with locally advanced or metastatic non-small-cell lung cancer (INFORM; C-TONG 0804): a multicentre, double-blind randomised phase 3 trial. **Lancet Oncol** **13**: 466-75, 2012.
 91. Cappuzzo F, Ciuleanu T, Stelmakh L, et al. Erlotinib as maintenance treatment in advanced non-small-cell lung cancer: a multicentre, randomised, placebo-controlled phase 3 study. **Lancet Oncol** **11**: 521-9, 2010.
 92. Miller VA, O'Connor P, Soh C, et al. A randomized, double-blind, placebo-controlled, phase IIIb

- trial (ATLAS) comparing bevacizumab (B) therapy with or without erlotinib (E) after completion of chemotherapy with B for first-line treatment of locally advanced, recurrent, or metastatic non-small cell lung cancer (NSCLC). **J Clin Oncol 27 (15 suppl):** abstract LBA8002, 2009.
93. Kabbinar FF, Miller VA, Johnson BE, et al. Overall survival (OS) in ATLAS, a phase IIIb trial comparing bevacizumab (B) therapy with or without erlotinib (E) after completion of chemotherapy (chemo) with B for first-line treatment of locally advanced, recurrent, or metastatic non-small cell lung cancer (NSCLC). **J Clin Oncol 28 (15 Suppl):** abstract 7526, 2010.
94. Perol M, Chouaid C, Perol D, et al. Randomized, phase III study of gemcitabine or erlotinib maintenance therapy versus observation, with predefined second-line treatment, after cisplatin gemcitabine induction chemotherapy in advanced non-small-cell lung cancer. **J Clin Oncol 30:** 3516-24, 2012.
95. Gerber DE, Schiller JH. Maintenance chemotherapy for advanced non-small-cell lung cancer: new life for an old idea. **J Clin Oncol 31:** 1009-20, 2013.
96. Cohen MH, Williams GA, Sridhara R, et al. FDA drug approval summary: gefitinib (ZD1839) (Iressa) tablets. **Oncologist 8:** 303-6, 2003.
97. Cohen MH, Williams GA, Sridhara R, et al. United States Food and Drug Administration drug approval summary: Gefitinib (ZD1839; Iressa) tablets. **Clin Cancer Res 10:** 1212-8, 2004.